

## Care & Maintenance

# Growing the Most Beautiful Lawn In the Neighborhood

## Part Three

# Care and Maintenance

### Irrigation for New Installation of Turfgrass Sod

Adequate moisture is necessary to promote optimum growth, density, color and durability of your turf grass. Water sod lightly after each 200 square feet is installed to prevent wilting or water shock. Once installation is complete, soak the entire area thoroughly. Keep the lawn soaked for the first week.

During extreme heat, water four to five times a day for short periods. In the second week, cut back to once or twice a day. Be sure the sod has ample moisture. Do not assume that if the soil under the sod is wet that the sod has enough moisture. Make your judgement call based on the moisture in the pad of the new sod.

After the first two weeks, roots have grown into the soil below and can pick up some of the moisture necessary to support the survival of the sod. Watering can now be reduced to encourage deeper root development into the




soil. Care must be taken to provide adequate water in accordance to heat, wind, humidity and sun as you begin to make the transition from frequent water to somewhat infrequent watering. This transition period will last an additional two to four weeks. As the roots develop deeper, the grass can go longer periods of time between water applications. Good wisdom and judgement must prevail as you make the transition from the water demands of newly installed sod to that of an established lawn.

The best lawns will result from deep and infrequent watering. The amount and frequency of watering depends on soil type, climate and topographic features.

Our other guide; Watering: When and How Much is a guide to help determine how to water your lawn as it matures.

### Fertilization

Fertilization applications are necessary for a healthy, dense, pest-free, durable and attractive lawn. In general, established lawn require nitrogen (N), phosphorous (P), potassium (K) and sulfur (S). Suggested formulas may be 20-10-10-5 or 25-10-10-13. Nitrogen promotes green color, tillering and leaf growth and is necessary for protein formation. Phosphorus is essential for cell development and root growth. Potassium promotes vigor, disease resistance and winter hardiness. Sulfur is involved with protein synthesis, growth and color. In certain situations, additions of some minor elements such as iron and zinc may be required.



Fertilizer should be applied three times a year. The first application should be applied in mid to late September and should supply 1/3 of the yearly nutrients. A second 1/3 application should follow after all moving has stopped; approximately very late October to early November, depending on the progression of fall weather into winter. It is important that this late application consists of water soluble nitrogen so that the root system can absorb it before winter. The last 1/3 of the annual fertilizer total is best applied in mid to late May and should contain nitrogen in a slow soluble form. To enhance and prolong good green color, the spring formulation should also contain some iron sulfate. If not, it can be supplied separately later during warmer weather to promote greening without additional increases in the foliar growth rate.





## Growing the Most Beautiful Lawn In the Neighborhood Part Three

# Care and Maintenance

### Fertilization continued...

These fertilizer recommendations are for medium to heavy textured soils. Application rates and intervals between applications need to be adjusted to match turf grass requirements. It is best to consult with experienced turf grass managers to obtain proper recommendations.

### Mowing

Proper mowing is necessary to maintain the appearance, health, vigor and density of the lawn. When to mow should be determined by the species and variety, growth rate and cutting height - not by the calendar. More frequent mowing is required during rapid growth periods in the spring and fall than during mid-summer.

Set the mower cutting bar by first placing the mower on a smooth surface, such as a sidewalk and measuring the bar height with a small ruler. Mow frequently enough that you remove only 1/3 of the grass at any one mowing. For example, to maintain your lawn at a 2 inch height, mow the grass when it reaches 3 inches. Closer mowing stresses the plants and depletes root reserves. Allowing turf grass to grow tall causes open, stemmy, poor density turf.



To maintain an attractive, well groomed lawn, keep mower blades very sharp at all times. Dull blades shred and bruise the leaves, causing light brown tips on the grass blades. Damaged grass is more susceptible to stress and diseases than healthy grass.

The return of grass clippings to the lawn, and eventually to the soil, have been considered to be a naturally accepted part of maintaining a lawn by the true turf experts. Grass clippings are a valuable resource. The clippings usually contain over 4 percent nitrogen, about 2 percent potassium and one-half percent phosphorus, as well as lesser amounts of other essential plant nutrients. By recycling the clippings the need for fertilizers and subsequent problems with surface and groundwater pollution are reduced.

Grass clippings are rapidly decomposed by bacteria and fungi, and do not contribute to thatch. Thatch results from the abnormally fast growth of tissues high in lignin, such as roots, rhizomes, stolons and crowns.

For most lawns, there is no need to remove the clippings from the lawn as long as they are small enough to work down into the grass. Clippings that are too long remain on the turf and can pack it down, damaging the growing plants. These clippings should be raked off to allow the grass to continue growing. Compost the clippings, provided that herbicides have not been applied to the lawn. Use the compost to fertilize the lawn and gardens.





## Care & Maintenance

# Growing the Most Beautiful Lawn In the Neighborhood Part Three

## Care and Maintenance

### Weed Control

Proper lawn maintenance practices are the most important weed control practices used on a home lawn. Thick, densely growing turf keeps weed problems to a minimum, and proper fertilizing, mowing, watering, aeration and thatch control are the first line of defense against weeds.

Herbicides, which kill existing weeds or prevent weeds from developing from seeds in the soil and turf, should be used after proper lawn maintenance practices have been established - not before. In this way, herbicides need be used only occasionally to treat weed problems.

To control broadleaf weeds, such as dandelion, plantain, or clover, selective post-emergent herbicides can be used. These kill certain plants (in this case broadleaf weeds) without damaging the grass plants. Apply them when the weeds are young and actively growing, taking care to prevent any herbicide drift onto desirable broad-leaved plants, such as flowers, shrubs, vegetables, or trees.



Pre-emergent herbicides generally provide the most effective control of annual grass weeds, such as crabgrass and foxtail. These herbicides kill weed seedlings just after the seed germinates, so the herbicides must be on the soil when the weed seeds germinate.



During a normal year, apply pre-emergent herbicides to control crabgrass by April 1st in warmer areas and May 1st in cooler areas. Our turf specialists can advise you on application times for your area. Do not wait until weeds have emerged to apply a pre-emergent herbicide; the best method is to apply the herbicide to areas of the lawn which had a crabgrass or foxtail problem in the previous year.

Perennial grass weeds, such as quackgrass, can be controlled chemically with non-selective, post-emergent herbicides. These materials can kill or damage all plants that they contact, including desirable grass plants. Consequently, they are only applied to weed hot spots. You have to re-seed spot treated areas.

**With proper care, you will have the best lawn in the neighborhood.**



2160 N Lake Hazel Rd, Meridian, ID 83642  
208-888-3760 800-488-8873(TURF)